PublisherInfo				
PublisherName		BioMed Central		
PublisherLocation		London		
PublisherImprintName	$\Box$	BioMed Central		

## Lively lipids

ArticleInfo		
ArticleID	:	4090
ArticleDOI	:	10.1186/gb-spotlight-20010521-01
ArticleCitationID	:	spotlight-20010521-01
ArticleSequenceNumber	:	161
ArticleCategory	:	Research news
ArticleFirstPage	:	1
ArticleLastPage	:	2
ArticleHistory	·	RegistrationDate : 2001–05–21 OnlineDate : 2001–05–21
ArticleCopyright	:	BioMed Central Ltd2001
ArticleGrants	:	
ArticleContext	:	130592211

## Jonathan B Weitzman

Email: jonathanweitzman@hotmail.com

In the May 18 Science, Farber *et al.* report the use of fluorescently quenched phospholipid substrates to screen for genes affecting lipid metabolism in living zebrafish larvae (*Science* 2001, **292:**1385-1388). The lipids were swallowed by the larvae and cleaved by lipases in the intestine before the fluorescent metabolites were transported to the liver and gall bladder. The assay demonstrated lipid metabolism defects in zebrafish mutants that have abnormal digestive organ morphology, as well as the '*fat free*' mutant that has an apparently normal digestive tract appearance. The authors suggest that the *fat free* mutation affects bile synthesis or secretion. This combination of mutagenesis screens with simple imaging of enzymatic function *in vivo* should provide insights into human lipid metabolism disorders.

## References

- 1. Science, [http://www.sciencemag.org]
- 2. Characterization of Ca2+-dependent phospholipase A2 activity during zebrafish embryogenesis.
- 3. Mutations affecting development of zebrafish digestive organs.

This PDF file was created after publication.